



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ARCTIC HORNED OWL.—One, nearly white, seen December 2 and 3. This is the only one observed in Wisconsin.

AMERICAN HAWK OWL.—One taken December 10. The first seen in in five years.

EVENING GROSBEEK.—Abundant in all suitable places.

AMERICAN CROSSBILL.—A flock of about twenty individuals as seen daily from December 2 to 10.

REDPOLL.—Several small flocks seen during the month.

SNOWFLAKE.—Three small flocks seen. Less than usual.

BOHEMIAN WAXWING.—About forty individuals visited the cedar trees in a neighbor's yard from November 28 to December 5.

NORTHERN SHRIKE.—Three seen. A regular winter visitant in small numbers.

RESIDENT.

BOB-WHITE.—Three small covies seen. Less than usual.

RUFFED GROUSE.—Five observed. Less than usual.

PRAIRIE HEN.—A covey of about thirty visited a corn field near by about every day. About as abundant as usual.

BARRED OWL.—One seen December 19.

GREAT HORNED OWL.—Eight observed.

HAIRY WOODPECKER.—Seen every day.

DOWNY WOODPECKER.—Four seen. Rare in winter, common in summer.

PILEATED WOODPECKER.—Three seen December 19. Regular resident in small numbers.

BLUE JAY.—Abundant everywhere.

AMERICAN GOLDFINCH.—One small flock observed.

BROWN CREEPER.—Several observed.

WHITE-BREASTED NUTHATCH.—Abundant in all woodlands.

CHICKADEE.—Abundant. Associates with White-breasted Nuthatch and Downy Woodpecker.

J. N. CLARK, *Meridian, Wis.*

GENERAL NOTES.

PHILADELPHIA VIREO, *Vireo philadelphicus*.—I shot a fine plumaged female about 7:15 A. M. September 19, 1898. It flew from the clearing side into a young hickory tree—which stood out in the sweet fern, detached from the woods, and disappeared in the foliage, soon coming in

view as it searched for food, flying from branch to branch and running out to the tips in a sprightly fashion without the warbler-like nervousness. Since John Cassin took the type specimen near Philadelphia in September, 1842, but six others have been recorded in eastern Pennsylvania. Eight captures in fifty-six years and all of them autumnal!

FRANK L. BURNS, *Berwyn, Penna.*

A PISCIVOROUS YELLOW-LEG.—While walking through the central market of this place, the other day, my attention was attracted by something protruding from the bill of a specimen of *Totanus melanoleucus*. Upon closer inspection this proved to be a minnow. The bird's œsophagus was literally jammed full of top minnows, (*Fundulus diaphanus*). When I made this discovery my mind at once wandered back to a certain railroad bridge which spans the Iowa river about two miles south of Iowa City, Ia., and the wintry scene changed to one of May. Here on the bridge I sat again and watched the trim form of Yellow-leg chasing minnows. Yes, I am convinced it was minnows he was after. Many times have I been seated here and watched this bird, which is quite common along the Iowa river during migration, wade out into the shallow water of the bars, moving along slowly with tilting gate, suddenly lower that long head and neck and proceed to run through the water at a speed which would have done credit to a college sprinter, quickly striking to right and left with his bill. Of course I thought some luckless water beetle had met its doom, but now I am convinced that minnows and not beetles caused the vicious movement.

P. BARTSCH, *Smithsonian Institute, Washington, D. C.*

LATE DATE FOR CATBIRD.—Two acquaintances, reputable men on whose word I can rely, observed a Catbird, *Galeoscoptes carolinensis*, feeding upon the pokeberries of which there has been an abundant crop, at the west side of the Valley creek dam, Valley Forge, on the 1st day of December, 1898. It was easily approached and frequently uttered the unmistakable *meew* as it busied itself in and out of the bushes as only a Catbird or Wren can do. In spite of the deep snow and freezing weather it appeared plump and fluffy, and as far as they could observe, was without any physical defect whatever. It was seen on five different occasions during the day.

FRANK L. BURNS, *Berwyn, Penna.*